arXiv:cond-mat/0201034v1 3 Jan 2002

Thermal Fluctuations of a Liquid Helium Inside a Spherical Volume

A.V.Kirilyuk[a] and A.V.Zvelindovsky[b]

[a] Department of Theoretical Physics, Odessa I.I.Mechnikov State University
Dvoryanskaya 2, 270026 Odessa, Ukraine
[b] Department of Biophysical Chemistry, University of Groningen
Nijenborgh 4, 9747 AG Groningen, The Netherlands

The theory of the thermal hydrodynamic fluctuations in bounded systems has been developed only for non-superfluid liquids on the base of the linearized Navier-Stokes equations [1]. However, the study of effect of confined geometry on correlation functions of superfluid liquid is important for explanation of experiments in which capillaries, pores, dense suspensions and other objects are probed by radiation [2].

In this paper we present the correlation theory of the equilibrium thermal hydrodynamic fluctuations of compressible superfluid helium confined by the spherical cavity of radius R. The hydrodynamic fluctuations in the superfluid helium are described on the base of linearized two-liquids Khalatnikov's equations [3] with spontaneous heat fluxes and strains. For mass flow $\delta\vec{J}$ and relative velocity $\delta\vec{w}=\vec{v}_n-\vec{v}_s$ of normal-fluid and superfluid components these equations have form

$$\frac{\partial\delta\rho}{\partial t}+\nabla\cdot\delta\vec{J}=0,\ \ \rho\frac{\partial\delta s}{\partial t}+\rho_s s\nabla\cdot\delta\vec{w}-\frac{\kappa}{T}\Delta\delta T=-\frac{1}{T}\nabla\cdot\vec{q},$$

$$\frac{\partial\delta\vec{J}}{\partial t}+\nabla\delta p-\frac{\eta}{\rho}\Delta\delta\vec{J}-\frac{1}{\rho}(\frac{1}{3}\eta+\zeta_2)\nabla(\nabla\cdot\delta\vec{J})-\frac{\rho_s\eta}{\rho}\Delta\delta\vec{w}+\frac{\rho_s}{\rho}(\rho\zeta_1-\frac{1}{3}\eta-\zeta_2)\nabla(\nabla\cdot\delta\vec{w})=-\vec{f},$$

$$\frac{\partial\delta\vec{J}}{\partial t}-\rho_n\frac{\partial\delta\vec{w}}{\partial t}+\nabla\delta p-\rho s\nabla\delta T-\zeta_1\nabla(\nabla\cdot\delta\vec{J})+\frac{\rho_s}{\rho}(-\rho\zeta_1+\rho^2\zeta_3)\nabla(\nabla\cdot\delta\vec{w})=-\rho\nabla H, \qquad (1)$$

where

$$\delta p=\left(\frac{\partial p}{\partial\rho}\right)_T\delta\rho+\left(\frac{\partial p}{\partial T}\right)_\rho\delta T,\quad \delta s=\left(\frac{\partial s}{\partial\rho}\right)_T\delta\rho+\left(\frac{\partial s}{\partial T}\right)_\rho\delta T.$$

Here s is the entropy per unit mass, p is the pressure; κ, η, and ζ_1,ζ_2,ζ_3 are the heat conductivity, the "first" viscosity, and the "second" viscosity coefficients, respectively; ρ_s,ρ_n, and ρ are the superfluid, normal-fluid, and total densities. The system (1) is written with account of the Onsager symmetry principle for the kinetic coefficients. All kinetic coefficients and thermodynamic derivatives in (1) assumed to be independent of coordinates. By (1) the fluctuation fields of mass flow, mass density and other variables are linear functionals of random forces in the right parts of these equations.

Assuming that the rigid-cavity surface is motionless we apply zero boundary conditions for the perpendicular component of mass flow $\delta\vec{J}$ and temperature fluctuations δT in the liquid helium, and the stick boundary conditions for the normal component of velocity [4]

$$\delta J_r(\vec{r},t)=0,\ \ \vec{v}_n(\vec{r},t)=0,\ \ \delta T(\vec{r},t)=0\ \ (r=R). \tag{2}$$

We present the mass flow, relative velocity, mass density and the temperature fluctuations in the form

$$\delta\vec{J}(\vec{r},t)=\sum_\lambda\left(J^L_\lambda(t)\vec{L}_\lambda(\vec{r})+J^M_\lambda(t)\vec{M}_\lambda(\vec{r})+J^N_\lambda(t)\vec{N}_\lambda(\vec{r})\right), \tag{3}$$

$$\delta\vec{w}(\vec{r},t)=\sum_\lambda\left(W^L_\lambda(t)\vec{L}_\lambda(\vec{r})+W^M_\lambda(t)\vec{M}_\lambda(\vec{r})+W^N_\lambda(t)\vec{N}_\lambda(\vec{r})\right), \tag{4}$$

$$\delta\rho(\vec{r},t)=\sum_\lambda R_\lambda(t)j_n(kr)Y_{mn}(\theta,\varphi), \tag{5}$$

$$\delta T(\vec{r},t)=\sum_\lambda T_\lambda(t)j_n(kr)Y_{mn}(\theta,\varphi), \tag{6}$$

where $\lambda=\{n,l,m\}$ and

$$\vec{L}_{mn}(\vec{r})=\frac{1}{k_\lambda}\nabla(Y_{mn}(\theta,\varphi)j_n(kr)),$$

$$\vec{M}_{mn}(\vec{r})=rot(\vec{r}Y_{mn}(\theta,\varphi)j_n(kr)),\quad \vec{N}_{mn}(\vec{r})=\frac{1}{k_\lambda}rot\,\vec{M}_{mn}(\vec{r}). \tag{7}$$

Here $Y_{mn}(\theta,\varphi)$ are the spherical functions and $j_n(k_\lambda r)$ are the spherical Bessel functions [5]. Note that $\vec{L}_{mn}(\vec{r})$, $\vec{M}_{mn}(\vec{r})$, $\vec{N}_{mn}(\vec{r})$ form the system of pairwise orthogonal functions.

From the boundary conditions (2) we find out eigenvalues k_λ:

$$j_n(\beta_{nl})=0,\quad k_\lambda=\beta_{nl}/R, \tag{8}$$

$$\gamma_{nl}j'_n(\gamma_{nl})+\alpha j_n(\gamma_{nl})=0,\quad \alpha=-n,n+1,\quad k_\lambda=\gamma_{nl}/R. \tag{9}$$

We present also the longitudinal and transverse parts of the random viscous force $\vec{f}$ and heat fluxes $\vec{q}$ in the form of expansion (3) with coefficients f^a_λ and Q^a_λ, respectively, $a=M,N,L$, and the random potential H - as in eq.(5) with coefficients H_λ.

It is convenient to use the Fourier representation for random $\delta\vec{J}$, $\delta\vec{w}$, $\delta\rho$, δT

$$x(t)=\int\limits_{-\infty}^{\infty}x_\omega exp[-i\omega t]d\omega. \tag{10}$$

Since the equations for the Fourier-amplitudes of the mass flow $J^b_{\lambda\omega}$ and relative velocity $W^b_{\lambda\omega}$ fluctuations ($b=M,N$) do not contain components of other fields we immediately obtain that

$$J^{M,N}_{\lambda\omega}=\rho_n W^{M,N}_{\lambda\omega}=\frac{f^{M,N}_{\lambda\omega}}{i\omega-k^2_\lambda\eta/\rho_n}\quad. \tag{11}$$

Ignoring the products of dissipation coefficients, fluctuation forces and keeping only the terms linear in the dissipation coefficients we get

$$D_\lambda J^L_{\lambda\omega} = -\frac{1}{T}\omega^2 k_\lambda^2 \rho_n \left(\frac{\partial p}{\partial T}\right)_\rho Q^L_{\lambda\omega} - i\omega k_\lambda^3 \rho_s \rho s \left(\frac{\partial p}{\partial T}\right)_\rho H_{\lambda\omega} +$$

$$+ i\omega \left(k_\lambda^2 \rho_s s \left(\frac{\partial p}{\partial T}\right)_\rho - k_\lambda^2 \rho_s \rho s^2 + \omega^2 \rho_n \rho \left(\frac{\partial s}{\partial T}\right)_\rho \right) f^L_{\lambda\omega}, \tag{12}$$

where

$$D_\lambda = -\rho_n \rho \left(\frac{\partial s}{\partial T}\right)_\rho \prod_{j=1}^{2} (\omega - k_\lambda u_j + i\alpha_{j\lambda} u_j)(\omega + k_\lambda u_j + i\alpha_{j\lambda} u_j) \tag{13}$$

and u_1, u_2 are the first- and second-sound velocities for the infinite space [3]; $\alpha_{1\lambda}$, $\alpha_{2\lambda}$ are the attenuation coefficients of the first- and second-sound, respectively. Ignoring terms proportional to $\rho_s \left(\frac{\partial \rho}{\partial T}\right)_p$ and $\frac{\kappa}{c_p}\left(\frac{c_p}{c_v} - 1\right)\frac{u_2^2}{u_1^2}$ for the first- and second-sound attenuation coefficients we obtain

$$\alpha_{1\lambda} = \frac{k_\lambda^2}{2\rho u_1} \left[\frac{4}{3}\eta + \zeta_2 + \frac{\kappa}{c_p}\left(\frac{c_p}{c_v} - 1\right) \right],$$

$$\alpha_{2\lambda} = \frac{k_\lambda^2}{2\rho u_2} \frac{\rho_s}{\rho_n} \left[\frac{4}{3}\eta + \zeta_2 - 2\rho\zeta_1 + \rho^2 \zeta_3 + \frac{\rho_n \kappa}{\rho_s c_p} \right], \tag{14}$$

where c_p, c_v are the specific heats at constant pressure and volume, respectively. The expansion coefficients for the excess mass density can be expressed in terms of J^L_λ.

From the expression for the rate of energy dissipation in the liquid HeII confined in the sphere volume V

$$Q(t) = \int_V dV \left(-\vec{v}_n \vec{f} + \rho_s \delta\vec{w}\, \nabla H - \frac{1}{T}\delta T\ \nabla \cdot \vec{q} \right) \tag{15}$$

we find that

$$Q_\omega = \sum_{a,\lambda} \left(-\frac{1}{\rho}(\rho_s W^a_{\lambda\omega} + J^a_{\lambda\omega})\Lambda^a_\lambda f^a_{\lambda\omega} + \frac{1}{3} k_\lambda \rho_s W^L_{\lambda\omega} \Lambda^L_\lambda H_{\lambda\omega} \right), \quad a = L, M, N, \tag{16}$$

where $\Lambda^a = \int_V d\vec{r}\, |\vec{a}(\vec{r})|^2$, $a = M, N, L$ are the normalization factors.

Using the Langevin equations (11), (12) we construct as in [6] the matrix of the generalized susceptibilities. And then by FDT find the spectral densities of equilibrium thermal fluctuations for the expansion amplitudes of the hydrodynamic fields. This resulted in

$$< J^L_\lambda J^{L\,*}_{\lambda'} >_\omega = \mathrm{Re}\, \frac{\Theta(\omega, T)\delta_{\lambda\lambda'}}{\Lambda^L_\lambda \pi} \frac{i\omega}{D_\lambda} \rho^2 \rho_n \left(\frac{\partial s}{\partial T}\right)_\rho (k_\lambda^2 u_2^2 - \omega^2),$$

$$< J^b_\lambda J^{b\,*}_{\lambda'} >_\omega = \mathrm{Re}\, \frac{\Theta(\omega, T)\rho\delta_{\lambda\lambda'}}{\Lambda^b_\lambda \pi(-i\omega + k_\lambda^2 \eta/\rho_n)}, \quad b = M, N, \tag{17}$$

where $\Theta(\omega, T)$ is the expression [6] for the average energy of the quantum oscillator.

The result differs from the spectral densities for an infinite superfluid by only the discrete behaviour of the wave numbers k_λ. Factorisation of the coefficient D_λ in (17) allows us to decompose the expressions for the pair correlation function of J^L_λ into the simple fractions. And the spectral densities of the J^L_λ as in the case of an infinite medium [7], consist of two contributions

$$< |J^L_\lambda|^2 >_\omega = \frac{2\rho\Theta(\omega,T)\omega^2(k_\lambda^2 u_2^2 - \omega^2)}{\pi\Lambda^L_\lambda k_\lambda^4 (u_1^2 - u_2^2)^2} \left\{ \frac{(\alpha_{1\lambda}u_1 - \alpha_{2\lambda}u_2)(\omega^2 - k_\lambda^2 u_1^2) - \alpha_{1\lambda}u_1 k_\lambda^2 (u_1^2 - u_2^2)}{[(\omega - k_\lambda u_1)^2 + \alpha_{1\lambda}^2 u_1^2][(\omega + k_\lambda u_1)^2 + \alpha_{1\lambda}^2 u_1^2]} + \right.$$

$$\left. + \frac{(\alpha_{2\lambda}u_2 - \alpha_{1\lambda}u_1)(\omega^2 - k_\lambda^2 u_2^2) + \alpha_{2\lambda}u_2 k_\lambda^2 (u_1^2 - u_2^2)}{[(\omega - k_\lambda u_2)^2 + \alpha_{2\lambda}^2 u_2^2][(\omega + k_\lambda u_2)^2 + \alpha_{2\lambda}^2 u_2^2]} \right\}, \quad k_\lambda = \gamma_{nl}/R. \tag{18}$$

Now it is easy to construct the spectral densities of thermal fluctuations of the mass flow and the mass density hydrodynamic fields

$$< \delta\vec{J}(\vec{r}, t)\delta\vec{J}(\vec{r}\,', t') >_\omega = \sum_{\lambda, A} < |J^A_\lambda|^2 >_\omega \vec{A}_\lambda(\vec{r})\vec{A}^*_\lambda(\vec{r}\,'), \;\; A = M, L, N, \tag{19}$$

$$< \delta\rho(\vec{r}, t)\delta\rho(\vec{r}\,', t') >_\omega = \sum_{\lambda} < |R_\lambda|^2 >_\omega j_n(k_\lambda r) j_n(k_\lambda r') Y_{mn}(\theta, \varphi) Y_{mn}(\theta', \varphi'), \tag{20}$$

where

$$< |R_\lambda|^2 >_\omega = \frac{k_\lambda^2}{\omega^2} < |J^L_\lambda|^2 >_\omega . \tag{21}$$

We have separated the first- and second-sound contributions to the mass density - mass density correlation function. The correlation functions are presented as the sum of the series in different spherical harmonics and roots of the transcendental equations (8), (9). As mentioned above, the difference between this case and that of infinite space is only in the discrete nature of the wave numbers k_λ. Each term in this series generates two Mandelstam-Brillouin doublets, which are caused by quasiphonons of two kinds, one having the first- and the other one the second-sound speed. Each components of the doublets are of Lorentz type, their widths being given by $\alpha_{1\lambda}u_1$ and $\alpha_{2\lambda}u_2$, and the spectral densities of the thermal fluctuations of mass flow and mass density can be presented as combinations of infinite numbers of such Lorentzians.

References

1. A.V. Zvelindovsky and A.V. Zatovsky, Nuovo Cimento D, 19(1997) 725.
2. D.L.Johnson, J.Phys.: Condens.Matter, 2(1990) SA449.
3. I.M.Khalatnikov, Introduction to the Theory of Superfluidity, Benjamin, New York, 1965.
4. L.D.Landau and E.M.Lifshitz, Fluid Mechanics, Pergamon, Oxford, 1982.
5. P.Morse and H.Feshbach, Methods of Theoretical Physics, Vol.2, McGraw-Hill, New York, 1953.
6. S.M.Rytov, Yu.A.Kravtzov, V.I.Tatarsky, Introduction to the Statistical Radiophysics, vol.2, Nauka, Moscow, 1978(in Russian).
7. Seth J. Putterman, Superfluid Hydrodynamics, North-Holland, Amsterdam, 1974.